

Algal Genomics

7A.1

NATURE, NURTURE AND THE STRUCTURE OF MACROALGAL GENOMES

J. Mark Cock^{1,2} (cock@sb-roscoff.fr), Jonas Coll  n^{1,2}, Lieven Sterck^{3,4}, Pierre Rouz  ^{3,4}, Delphine Scornet^{1,2}, Veronique Anthouard⁵, Fran  ois Artiguenave⁵, Jean-Marc Aury⁵, Kenny Billiau^{3,4}, Eric Bonnet^{3,4}, John H.F. Bothwell^{6,7,8}, Loraine Brillet⁹, Wilfrid Carre⁹, Susana M. Coelho^{1,2}, Erwan Corre⁹, Corinne Da Silva⁵, Claire Jubin⁵, Cindy Martens^{3,4}, Florian Maumus¹⁰, Diego Miranda-Saavedra¹¹, Akira F. Peters^{1,2}, Betina Porcel⁵, Hadi Quesneville¹², Catherine Boyen^{1,2}, Yves Van de Peer^{3,4} and Patrick Wincker⁵

¹UPMC Univ. Paris 06, The Marine Plants and Biomolecules Laboratory, UMR 7139, Station Biologique de Roscoff, Place Georges Teissier, BP74, 29682 Roscoff Cedex, France; ²CNRS, UMR 7139, Laboratoire International Associ   Dispersal and Adaptation in Marine Species, Station Biologique de Roscoff, Place Georges Teissier, BP74, 29682 Roscoff Cedex, France; ³Department of Plant Systems Biology, VIB, B-9052 Ghent, Belgium; ⁴Department of Plant Biotechnology and Genetics, Ghent University, B-9052 Ghent, Belgium; ⁵CEA, DSV, Institut de G  nomique, G  noscope, 2 rue Gaston Cr  mieux, CP5706, 91057 Evry, France; ⁶Queen's University Belfast, School of Biological Sciences, 97 Lisburn Road, Belfast, BT9 7BL, UK; ⁷Queen's University Marine Laboratory, Portaferry, Co. Down, BT22 1PF, UK; ⁸Marine Biological Association of the UK, The Laboratory, Citadel Hill, Plymouth, PL1 2PB, UK; ⁹Computer and Genomics Resource Centre, FR 2424, Station Biologique de Roscoff, Place Georges Teissier, BP74, 29682 Roscoff Cedex, France; ¹⁰Institut de Biologie de l'Ecole Normale Sup  rieure (IBENS), Centre National de la Recherche Scientifique UMR8197, Ecole Normale Sup  rieure, 75005 Paris, France; ¹¹Department of Haematology, Cambridge Institute for Medical Research, University of Cambridge, Hills Road, Cambridge CB2 0XY, UK and ¹²Unit   de Recherches en G  nomique-Info (UR INRA 1164), INRA, Centre de recherche de Versailles, bat.18, RD10, Route de Saint Cyr, 78026 Versailles Cedex, France

The recent acquisition and analysis of complete genome sequences for two macroalgal species has provided unparalleled insights into the biology and evolutionary history of these important organisms. The recently published genome sequence of the filamentous brown alga *Ectocarpus siliculosus* (Cock *et al.*, Nature 2010) has not only provided clues as to how brown alga have adapted to the harsh conditions of the intertidal, but has also allowed the generation of hypotheses about the molecular events underlying the evolution of complex multicellular development in this group of organisms. The more recently completed genome sequence of the red alga *Chondrus crispus* (Coll  n *et al.*, unpublished) has similarly provided important information about the origins and the biology of the red algae. Both *Ectocarpus* and *Chondrus* are found in the intertidal zone, but they have had very different evolutionary histories and have been evolving independently for more than a billion years. Interestingly, they also exhibit very marked differences in terms of genome structure and organisation, suggesting that their evolutionary histories have played a more important role in shaping their genomes than their adaptation to the intertidal environment.

7A.2

DIVERSITY OF MARINE PRASINOPHYCEAE AND ASSOCIATED VIRUSES

Nigel Grimsley (nigel.grimsley@obs-banyuls.fr)

Oceanological Observatory of Banyuls, UMR 7232 CNRS – UPMC, 1 avenue de Fontaule, 66650 Banyuls sur mer, France

The order Mamiellales (Prasinophyceae) comprises freshwater and marine “green” unicellular photosynthetic eukaryotes. Many of these have very small cell sizes and are named ‘picoeukaryotes’ (cell diameter < 2 µm). The ecological importance of marine Mamiellales in coastal waters, their easy manipulation and their small genome sizes have fostered genome projects in several different species. In this presentation, recent analyses of the 6 available genomes from the *Bathycoccus*, *Micromonas* and *Ostreococcus* genera are presented and the insights gained into the species